

Black holes, cosmological singularities and change of signature

Marcus Kriele

Fachbereich Mathematik, Sekr. MA 8-3

Technische Universität Berlin

Strasse des 17 Juni 136, 10623 Berlin

Federal Republic of Germany

Jérôme Martin*

Department of Applied Mathematics and Theoretical Physics

University of Cambridge, Silver street

Cambridge CB3 9EW, United Kingdom

February 5, 2008

Abstract

There exists a widespread belief that signature type change could be used to avoid spacetime singularities. We show that signature change cannot be utilised to this end unless the Einstein equation is abandoned at the surface of signature type change. We also discuss how to solve the initial value problem and show to which extent smooth and discontinuous signature changing solutions are equivalent.

* Permanent adress: Laboratoire de Gravitation et Cosmologie Relativistes, Université Pierre et Marie Curie, CNRS/URA 769, Tour 22/12, Boite courrier 142, 4 place Jussieu 75252 Paris cedex 05, France.

1 Introduction

According to the Hawking-Penrose theorem [1], singularities in General Relativity seem to be unavoidable. The two most well-known examples are the singularities of the Friedmann-Robertson-Walker metric ("big bang singularities") and of the Schwarzschild solution ("black hole singularities"). The presence of these singularities is usually interpreted as the sign that General Relativity, a classical theory, is no more valid since quantum effects have to be taken into account when the curvature reaches the Planck limit. Therefore, it is not surprising that a possible solution to this problem has been suggested in the context of quantum cosmology. Recent studies [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12] have shown that change of signature can be also a feature of classical General Relativity. In this framework, the very early Universe is described by a Riemannian¹ manifold which does not have a big bang singularity [13]. It has been hoped that this is a consequence of signature change (this has been argued in [3] since the singularity theorems do not apply for Riemannian manifolds). However, in this paper we will show that big bang singularities which would occur without signature change will reappear as Big Crunch singularities. We also answer the question whether one can employ signature change in order to avoid black hole singularities to the negative.

There have been put forward different suggestions as to how to implement signature change classically. As a consequence, there are now different competing theories and an ongoing discussion about the relative merit of the smooth and discontinuous description of signature type change (cf. [2, 3, 4, 5, 12, 14]). These different proposals can be divided into two groups.

- (a) One imposes regularity conditions at the hypersurface of signature change which can be understood as imposing the Einstein equations (in a suitable form) at the surface of signature type change. This approach has been adopted by [2, 4, 5, 6, 7, 8, 9, 10, 12].
- (b) One views spacetime as a 1-parameter family of Riemannian manifolds and therefore relaxes the regularity conditions at the surface of signature change. This approach has been adopted by [3, 14].

We are of the opinion that wherever one can use the Einstein equation one should impose it and are therefore favouring approach (a) which we will consider exclusively. Within

¹In the physics literature the term 'Euclidean' is often used instead of the term 'Riemannian' which is more common in the mathematical literature.

approach (a) there are two competing proposals: One can implement signature change with a discontinuous but non-degenerate or with a continuous but degenerate metric. So naturally the question arises whether any of these two implementations is superior. In [12] this question has been attempted to decide from a geometrical viewpoint. The authors concluded that the smooth description was vastly superior. However, this conclusion rests on “a priori” demands on the regularity of the solutions. Here we show that for solutions of Einstein’s equations different regularity conditions arise naturally in the discontinuous description. Assuming these regularity conditions the space of solutions of the Einstein equations in either scenario are canonically equivalent. Thus it appears to be a matter of taste which setting one prefers.

Our paper is organized as follows: In section 2 we discuss the initial value problem and show the equivalence between continuous and discontinuous change of signature for a specific class of solutions. In section 3.1 we show that big bang singularities of Lorentzian solutions will reappear as Big Crunch singularities in the corresponding type changing solution. In section 3.2 we prove the impossibility of matching a Riemannian manifold inside the black hole horizon without introducing new singularities.

2 Comparison between continuous and discontinuous change of signature

Let us first recall the two definitions. In the discontinuous case one should restrict to signature change at *spacelike* hypersurfaces $\tilde{\mathcal{D}}$ because only in this case it is possible for $\tilde{\mathcal{D}}$ to inherit the same structure from both the Lorentzian and the Riemannian region. But given a distinguished spacelike hypersurface one can define a distinguished time function, the parameter function of the unit geodesics starting orthogonal to this hypersurface. To employ this natural time function has several advantages. For instance, it makes it possible to write down the energy momentum tensor as a well defined object and it facilitates the comparison of smooth and discontinuous signature change. We do not want to specify the regularity of the considered type changing metric yet. So let $\mathcal{G}$ be a subset of all functions $f : M \rightarrow \mathbb{R}$ to be specified later and define:

Definition 1 $(\tilde{M}, \tilde{g})$ is a *type changing spacetime with jump of class $\mathcal{G}$* if $\tilde{M}$ is a smooth, 4-dimensional manifold with an everywhere non-degenerate, symmetric $(0, 2)$ tensor field

$\tilde{g}$ which is continuous everywhere except at a hypersurface $\tilde{\mathcal{D}}$. For any x in $\tilde{\mathcal{D}}$ there exists a coordinate neighbourhood such that $\tilde{g}$ is given by $\tilde{g} = -\eta d\tilde{t}^2 + \tilde{g}_{ij}(\tilde{t}, \tilde{x}^1, \tilde{x}^2, \tilde{x}^3) d\tilde{x}^i d\tilde{x}^j$, where $\eta = \text{sign}(\tilde{t})$ and $\tilde{g}_{ij} \in \mathcal{G}$ ($i, j \in \{1, 2, 3\}$).

In [12] the class $\mathcal{G}$ has been taken as the class of C^k -functions. This has been justified since the $\tilde{t}$ -coordinate is the invariantly defined time function so that one can view the $\tilde{g}_{ij}$ as a C^k -differentiable 1-parameter family of C^k -3-metrics. However, this choice of class is not natural from a physical point of view as we will see below.

For the smooth case one may define [6]:

Definition 2 (M, g) is a *transverse, type changing C^k -spacetime* ($k \in \mathbb{N} \cup \{0, \infty, \omega\}$)² if M is a smooth, 4-dimensional manifold with a symmetric C^k -(0, 2)-tensor field g such that at any point $x \in \mathcal{D} := \{\hat{x} \in M \mid g|_{\hat{x}} \text{ is degenerate}\}$ we have $d(\det(g_{ab}))|_x \neq 0$ for some (and hence any) coordinate system. Moreover, at any point where g is non-degenerate it is either Riemannian or Lorentzian.

The main difference to definition 1 is that here g is assumed to be a smooth tensor field. It is clear that the surface of signature change must be given by $\det(g_{ab}) = 0$. The additional condition, $d(\det(g_{ab}))|_x \neq 0$, may be thought of as a genericity condition.

This definition also allows to have signature change at null surfaces. However, for cosmological applications one would like to have a spacelike surface of signature change. Thus we define

Definition 3 (M, g) has a *transverse radical* if in addition $\text{Rad}_x := \{v_x \in T_x M \mid g(v_x, \cdot) = 0\}$ intersects $T_x \mathcal{D}$ transversely for all $x \in \mathcal{D}$.

Observe that Rad_x is necessarily one-dimensional. Notice that since g is a well defined tensor field it is natural to consider C^k metrics g . In contrast to the discontinuous case we do not need to specify an (invariantly defined) system of coordinates. We will see below that the class of such metrics is also natural from a physical point of view.

Although these definitions are rather different in spirit one can introduce coordinates (t, x^1, x^2, x^3) such that the content of definitions 2, 3 can be reformulated similarly to definition 1:

²A function f is said to be C^ω if f is real analytic.

Lemma 4 *Let M be a manifold, g be a $(0,2)$ -tensor field, and $\mathcal{D} := \{\hat{x} \in M \mid g_{\hat{x}} \text{ is degenerate}\}$. Then (M, g) is a transverse, type changing spacetime with transverse radical if and only if for any $\hat{x} \in \mathcal{D}$ there exists a neighbourhood and coordinates (t, x^1, x^2, x^3) such that*

$$g = -tdt^2 + g_{ij}(t, x^1, x^2, x^3)dx^i dx^j.$$

Proof: This has been shown (in a more general context) in [8]. ■

It is now possible to relate the smooth picture to the discontinuous one. A necessary condition for equivalence is clearly that the Lorentzian and the Riemannian parts of the two descriptions are isometric. Thus two metrics $g, \tilde{g}$ are *equivalent* if there exists a homeomorphism which is an isometry away from the surfaces of signature type change. Since the surfaces $\tilde{t} = \text{const}$ and $t = \text{const}$ have an invariant meaning such a transformation must be given by

$$\psi: (t, x^1, x^2, x^3) \mapsto (\tilde{t}, \tilde{x}^1, \tilde{x}^2, \tilde{x}^3) \text{ with } \tilde{t} = \eta \frac{2}{3} (\eta t)^{3/2}, \quad \tilde{x}^i = \tilde{x}^i(\tilde{x}^1, \tilde{x}^2, \tilde{x}^3).$$

It is clear that we only need to consider transformations ψ with $\tilde{x}^i = x^i$.

Then $(M, \tilde{g})$ is equivalent to a metric transverse, type changing C^k -spacetime if and only if $(M, \psi^* \tilde{g})$ satisfies definition 2.

We will now consider the Einstein equations and show that the solutions in both the discontinuous and the smooth approach are canonically equivalent. Assume for definiteness³ that the energy momentum tensor has the form of a scalar field with arbitrary potential (but which is not coupled to the scalar curvature):

$$T_{ab} = \partial\phi_a \partial\phi_b - \frac{1}{2} (g(\text{grad}(\phi), \text{grad}(\phi)) + V(\phi)) g_{ab},$$

where V is some analytic function and ϕ is the scalar field. For convenience we will also assume that $M \setminus \mathcal{D}$ consists of two connected components, a Riemannian component $\tilde{M}^+$ and a Lorentzian component $\tilde{M}^-$.

In the discontinuous case (and arbitrary coordinates) the Einstein equations will in general fail to be distributional. Even in our adapted (shift = 0) coordinates $(\tilde{t}, \tilde{x}^1, \tilde{x}^2, \tilde{x}^3)$ the component of the Ricci tensor will in general fail to be defined distributionally. However, the energy momentum tensor is a well defined distributional tensor and it makes

³But compare the conclusion in which we point out a key feature of the scalar field energy momentum tensor which makes the following discussion possible

sense to demand that it is bounded. This requirement just means that at the surface of signature type change we do not have a singularity due to concentration of matter. From this condition it follows that $\partial_{\tilde{t}}g_{ij} = 0$ at $\tilde{t} = 0$ and that *then* also the components of the Ricci tensor are defined distributionally⁴.

The initial value problem at the surface of signature change splits into qualitatively different initial value problems, one for the Lorentzian and the other one for the Riemannian part.

Let us first consider the Lorentzian part. We will denote all quantities with a hat in order to distinguish from the signature changing case. The condition that $\hat{T}_{ab}$ is bounded was equivalent to the requirement $\partial_{\hat{t}}\hat{g}_{ij} = 0$ for the initial 3-metric. We therefore have only to solve the usual Einstein equation for this sort of initial data. This is (given smooth data) always possible if the usual constraint equation holds at the surface. The proof of Lemma 2 in [7] implies in addition that the Taylor series of $\hat{g}_{ij}$ and of the scalar field $\hat{\phi}$ depend smoothly on $\hat{t}^2$. Observe that in the analytic⁵ case at $\hat{t} = 0$ we not only have a surface of infinitesimal time symmetry but that the map $\hat{t} \rightarrow -\hat{t}$ induces an isometry.

For the Riemannian part it would be much more difficult to solve the initial value problem because the system of differential equations is not hyperbolic in this region. But one can construct a Riemannian solution from a Lorentzian solution employing the trick with the Wick rotation: If one replaces $\hat{t}$ with $\tilde{t} = i\hat{t}$ in the Lorentzian solution one automatically obtains a solution of the Riemannian equations. Moreover, this Riemannian solution has a real Taylor series and therefore is real if it is analytic (cf. appendix A for an illustration that one *must* impose analyticity).

It is now clear how to obtain a signature changing solution $(M, \tilde{g})$: We use the Wick rotation in one connected component of $M \setminus \mathcal{D}$ which we call $\tilde{M}^+$. Observe that the metric coefficients now depend analytically on $\eta\tilde{t}^2$. Thus $\mathcal{G}$ should be assumed to be the class of functions which depend analytically on $\eta\tilde{t}^2, \tilde{x}^1, \tilde{x}^2, \tilde{x}^3$.


In order to obtain a smooth signature type changing solution we now apply the transformation ψ to the discontinuous solution. This is possible since the discontinuous solution depends analytically on $\eta\tilde{t}^2$. This transformation results in an analytic solution which depends analytically on t^3 . Moreover, any analytic, transverse, type changing solution with

⁴The above follows immediately from the expressions in the appendix of [12]. Observe that a similar but less convoluted claim on the bottom of page 2366 in this paper is not true (this, however, does not affect the rest of the paper)

⁵I.e., each metric components $\hat{g}_{ij}$ can be expressed as a power series in the variables $(\hat{t}, \hat{x}^1, \hat{x}^2, \hat{x}^3)$

bounded energy momentum tensor can be obtained in this way [7].

We have therefore established the following diagram:



Example 1 In order to illustrate the previous diagram, let us consider the Friedmann-Robertson-Walker (FRW) metric with a cosmological constant (this example has been already studied in [2, 3]). In the case where the continuous choice is made, the metric can be written as:

$$ds^2 = \begin{cases} -tdt^2 + \cos^2(\frac{2}{3}(-t)^{\frac{3}{2}})d\Omega_3^2 & -(\frac{3\pi}{4})^{\frac{2}{3}} \leq t \leq 0 \\ -tdt^2 + \cosh^2(\frac{2}{3}t^{\frac{3}{2}})d\Omega_3^2 & t \geq 0 \end{cases}$$

where $d\Omega_3^2$ represents the line element of a three-sphere. The metric is manifestly degenerate at the surface $t = 0$ but is C^∞ since the Taylor series of $\cos(\frac{2}{3}(-t)^{\frac{3}{2}})$ and $\cosh(\frac{2}{3}t^{\frac{3}{2}})$ are the same:

$$\cos(\frac{2}{3}(-t)^{\frac{3}{2}}) = \cosh(\frac{2}{3}t^{\frac{3}{2}}) = 1 + \frac{1}{2!} \left(\frac{2}{3}\right)^2 t^3 + \frac{1}{4!} \left(\frac{2}{3}\right)^4 t^6 + \dots$$

From this expression, it is obvious that g_{ij} depends analytically on t^3 .

We can now perform the transformation $\tilde{t} = \frac{2}{3}\eta(\eta t)^{\frac{3}{2}}$, $\tilde{x}_i = x_i$; then the metric takes the form:

$$d\tilde{s}^2 = \begin{cases} d\tilde{t}^2 + \cos^2(\tilde{t})d\Omega_3^2 & -(\frac{\pi}{2}) \leq \tilde{t} \leq 0 \\ -d\tilde{t}^2 + \cosh^2(\tilde{t})d\Omega_3^2 & \tilde{t} \geq 0 \end{cases}$$

The metric is no more degenerate on the surface $\tilde{t} = 0$ but is discontinuous. $\tilde{M}^+$ is half of the sphere S^4 whereas $\tilde{M}^-$ is half of the De Sitter spacetime. $\tilde{g}_{ij}$ depends analytically on $\eta\tilde{t}^2$.

If we use the Wick rotation $\tilde{t} = i\hat{t}$ on $\tilde{M}^+$, we obtain the following metric on M :

$$ds^2 = -d\hat{t}^2 + \cosh^2(\hat{t})d\Omega_3^2.$$

M is now the entire De Sitter spacetime.

Finally, if we perform a Wick rotation on $(M, \hat{g})$, the metric becomes:

$$ds^2 = d\check{t}^2 + \cos^2(\check{t})d\Omega_3^2,$$

namely the metric of the sphere S^4 . Then, all the relationships of the diagram have been explicitly shown using the simple example of the FRW metric.

3 Spacetime singularities

3.1 Signature change in cosmology

We show that one cannot use signature type change in order to turn singular, inextendible spacetimes into non-singular ones:

Proposition 5 *Let (M, g) be a signature type changing spacetime. If it is singularity-free then the Lorentzian spacetime $(M, \hat{g})$ corresponding to it is singularity free.*

Proof: The discussion in section 2 shows that the Lorentzian spacetime $(M, \hat{g})$ corresponding to (M, g) admits a reflectional isometry $\hat{t} \mapsto -\hat{t}$. Hence if it has a singularity in $\tilde{M}^+$ then it also must have a singularity in $\tilde{M}^-$. Since $\tilde{M}^-$ is isometric to one connected component of $M \setminus \mathcal{D}$ in the Lorentzian solution our claim follows. ■

Actually, there is also a different argument first pointed out by Hayward [15]. Since the surface of signature change is totally geodesic in a closed cosmology the universe is immediately contracting unless one violates the strong energy condition. Hayward interpreted this circumstance as positive evidence for an inflationary phase in which the strong energy condition would be violated.

Notice that in example 1 above all solutions are singularity-free.

3.2 Signature change at black holes

One may speculate that it could be possible to avoid black hole singularities by imposing change of signature at the boundary of the black hole. In this section we show that a signature changing mechanism would not work. Recall that there are two possible definitions as to what the boundary of a black hole is:

- (a) From a quasi-local point of view it is natural to consider the outer trapping horizon as the boundary of the black hole [16].
- (b) One may also take the event horizon of a black hole as a definition of its boundary. However, this definition is essentially global and therefore does not capture the physical content as well as the definition in (a).

An outer trapping horizon is a hypersurface surface which is foliated by outer marginally trapped surfaces (outer means that the expansion of the null direction which vanishes would become negative when the surface is moved into the other null direction). Hayward has also shown that the outer trapping horizon is spacelike if the null energy condition and a genericity condition hold [16, Theorem 2]. Thus signature change at the trapping horizon would be mathematically equivalent to cosmological signature change. In particular, if the energy momentum tensor is supposed to be bounded then one has to assume that this surface is a surface of (infinitesimal) time symmetry. This is certainly impossible where a physical black hole develops. Thus one cannot implement signature type change at the trapping horizon.

It follows that signature change could only be implemented at the event horizon of a black hole. Observe that the event horizon is a null surface $\mathcal{D}$ and that the weakest regularity condition to impose on $\mathcal{D}$ is that the induced metric is unambiguously defined (ie. the metric inherited from the Lorentzian part should be the same as the metric inherited from the Riemannian part). Since in Riemannian geometry there do not exist non-vanishing null vectors it is clear that discontinuous signature type change is impossible.

However, continuous signature type change is not ruled out yet. As a trivial example consider the 2-dimensional metric $ydx^2 + dy^2$. This is a transverse, type changing metric for which the surface of signature change $\mathcal{D}$ is given by $y = 0$. However, the radical, $\text{span}\{\partial_x\}$ is tangent to $\mathcal{D}$ and hence it fails to be transverse. Thus $\mathcal{D}$ is a null surface. But if one calculates the Gauß curvature K one obtains $K = 1/y^2$ which diverges at $y = 0$

We will now show that signature change at a null hypersurface implies the existence of curvature singularities. If $\mathcal{D}$ is null then the radical Rad must be everywhere tangent to $\mathcal{D}$. Thus it suffices to prove the following proposition which is an extension of [17, Theorem 3]:

Proposition 6 *If (M, g) is a transverse type changing spacetime but the radical not transverse at $x \in \mathcal{D}$ then the energy momentum tensor T_{ab} is unbounded at x .*

Proof: There exists an adapted orthonormal frame $\{e_0, e_1, e_2, e_3\}$ such that $g(e_i, e_j) = \delta_j^i$, $g(e_0, e_i) = 0$ for $i, j \in \{1, 2, 3\}$ and $g(e_0, e_0) = \tau$ where τ is a function with $\tau|_{\mathcal{D}} = 0$ but $d\tau_x \neq 0$. The existence of such a frame follows with a slight adaptation of the Gram-Schmidt procedure [18]. Since $g(\nabla_{e_a} e_b, e_c)$ extends smoothly and since

$$\begin{aligned} g(\nabla_{e_a} \nabla_{e_b} e_a, e_b) &= \nabla_{e_a} g(\nabla_{e_b} e_a, e_b) - g(\nabla_{e_b} e_a, \nabla_{e_a} e_b) \\ &= -\frac{1}{\tau} g(\nabla_{e_b} e_a, e_0) g(e_0, \nabla_{e_a} e_b) + \text{smooth terms}, \end{aligned}$$

we get

$$\begin{aligned} g(R(e_a, e_b)e_a, e_b) &= -\frac{1}{\tau} \left(g(\nabla_{e_b} e_a, e_0) g(\nabla_{e_a} e_b, e_0) - g(\nabla_{e_a} e_a, e_0) g(\nabla_{e_b} e_b, e_0) \right) \\ &\quad + \text{smooth terms}. \end{aligned}$$

Setting $e_b = e_0$, observing that at x we have $0 = d\tau(e_0) = 2g(\nabla_{e_0} e_0, e_0)$, and using that $g([e_a, e_0], e_0)$ vanishes at $\mathcal{D}$, we obtain

$$g(R(e_a, e_0)e_a, e_0) = -\frac{1}{\tau} \left(g(\nabla_{e_a} e_0, e_0) \right)^2 + \text{smooth terms}.$$

Now notice that the frame $\{e_0, e_1, e_2, e_3\}$ can be chosen so that e_2, e_3 are tangent to $\mathcal{D}$. Thus at x we have $d\tau(e_0) = d\tau(e_2) = d\tau(e_3) = 0$ but $d\tau(e_1) \neq 0$. Since $g(\nabla_{e_a} e_0, e_0) = \frac{1}{2}d\tau(e_a)$ we conclude that $g(R(e_1, e_0)e_1, e_0)$ diverges (such that $\tau g(R(e_1, e_0)e_1, e_0)$ is bounded) whereas $g(R(e_a, e_0)e_a, e_0)$ is bounded for $a = 0, 2, 3$. In particular, $\text{Ric}(e_0, e_0)$ diverges such that $\tau \text{Ric}(e_0, e_0)$ and $(\text{Ric}(e_0, e_0) - g(R(e_1, e_0)e_1, e_0))/\tau$ are bounded. For $\alpha \in \{2, 3\}$ we have

$$\text{Ric}(e_\alpha, e_\alpha) = \sum_{b=0}^3 (g(e_b, e_b))^{-1} g(R(e_\alpha, e_b)e_\alpha, e_b).$$

$\tau \text{Ric}(e_\alpha, e_\alpha)$ is bounded since $g(R(e_\alpha, e_0)e_\alpha, e_0)$, $g(R(e_\alpha, e_b)e_\alpha, e_b)$, $\tau g(R(e_\alpha, e_b)e_\alpha, e_b)$, $(g(e_b, e_b))^{-1}$ are bounded for $b \neq 0$.

$$\begin{aligned} \text{Ric}(e_1, e_1) &= \sum_{b=0}^3 (g(e_b, e_b))^{-1} g(R(e_1, e_b)e_1, e_b) \\ &= \sum_{\alpha=2}^3 (g(e_\alpha, e_\alpha))^{-1} g(R(e_1, e_\alpha)e_1, e_\alpha) + \frac{1}{\tau} (\text{Ric}(e_0, e_0) + \text{smooth terms}). \end{aligned}$$

Thus we obtain for the scalar curvature $s = \frac{2}{\tau} (\text{Ric}(e_0, e_0) + \text{smooth terms})$. In particular it follows that $T(e_2, e_2) = -\frac{1}{\tau} \text{Ric}(e_0, e_0) + \frac{1}{\tau} (\text{smooth terms})$ diverges at a similar rate as $\frac{1}{\tau^2}$. ■

Observe that the singularity provided by proposition 6 is non-distributional since $\tau^2 T(e_2, e_2)$ does not converge to zero. This would be different if $\tau T(e_2, e_2)$ was bounded.

4 Conclusion

In section 2 we have shown that there exists a well defined equivalence of solutions of Einsteins equations for both the continuous and the discontinuous implementation of signature change. In order to establish this equivalence we have used an energy momentum tensor for a scalar field. The form of this energy momentum tensor is crucial for the argument. The key feature is that time derivatives occur only quadratically $(\partial_{\tilde{t}} a \partial_{\tilde{t}} b)$ and that each such quadratic pair is weighted with the $\tilde{t}\tilde{t}$ -component of the inverse of the metric. This circumstance is responsible for the fact that the Taylor series of the $\tilde{g}_{ij}$ depend quadratically on $\tilde{t}$. If this was not the case the Wick transformation could not give rise to real solutions. We doubt wether there exist any solutions for energy momentum tensors which do not have this property. In principle, it is possible to construct Lagrangians whose energy momentum tensor violates this property. For instance, one may have as matter quantities a scalar field ν and a vector field X and add a term $d\nu(X)$ to the Lagrangian.

We have established that it is a matter of taste whether one prefers continuous or discontinuous signature change, provided Einstein's equations are imposed. If one works in the discontinuous picture one has the advantage that the metric is nowhere degenerate and that there exist observer fields which can be smoothly continued into the Riemannian region. However, one has to pay the price that solutions of Einstein's equation have unusual regularity properties. For instance, due to the factor η in general the energy momentum tensor fails to be analytic unless spacetime is static [12]. On the other hand, observe that the transformation ψ transforms any such solution into an analytic transverse type changing solution for the analogous problem with an analytic energy momentum tensor. Thus, from a mathematical point of view, the continuous picture seems to be more familiar.

Finally, we have shown that signature change cannot be sucessfully employed in order to avoid singularities of solutions of Einstein's equations.

A Appendix

Consider a smooth but non-analytic Lorentzian solution which depends on $\tilde{t}^2$. Applying the Wick rotation we obtain a Riemannian solution of the corresponding Riemannian Einstein equations. This Riemannian solution has a real Taylor series. Nevertheless, this solution fails to be real in general. In order to illustrate this point consider the simple but completely analogous massless wave equation on a flat, 2-dimensional background. In the Lorentzian region it is the usual wave equation $-\partial_{\tilde{t}}\partial_{\tilde{t}}\phi + \partial_{\tilde{x}}\partial_{\tilde{x}}\phi = 0$ whereas in the Riemannian part it is the usual Laplace equation $\partial_{\tilde{t}}\partial_{\tilde{t}}\phi + \partial_{\tilde{x}}\partial_{\tilde{x}}\phi = 0$, both with initial condition $\partial_{\tilde{t}}\phi(0, \tilde{x}) = 0, \phi(0, \tilde{x}) = \vartheta(\tilde{x})$ for some arbitrary function ϑ . Although for any smooth initial function ϑ the Lorentzian region has a unique, smooth solution $\phi(\tilde{t}, \tilde{x}) = \vartheta(\tilde{x} + \tilde{t}) + \vartheta(\tilde{x} - \tilde{t})$ it is well known that any smooth solution in the Riemannian region must have *analytic* initial data [19, p. 455]. Of course, we could construct a solution of the Riemannian equation using Wick rotation. However, this solution fails to be real even though its Taylor series is automatically real. As an explicit example, consider the function

$$\vartheta(x) := \begin{cases} 0 & \text{for } x \geq 0 \\ xe^{-1/x^2} & \text{for } x < 0. \end{cases}$$

The Taylor series of this function centred at $\tilde{t} = 0$ vanishes and therefore is real. But it is clear that $\phi(\tilde{t}, \tilde{x})$ is not a real valued function.

Thus one has to demand that the initial data are analytic.

Acknowledgement M. K. would like to thank the Laboratoire de gravitation et cosmologie relativiste of the Université Pierre et Marie Curie for warm hospitality. J. M. would like to thank the Ministère de la Recherche et de l'Enseignement Supérieur for a research grant. We would like to thank Robin Tucker for pointing out that to refer to ψ as a ‘coordinate transformation’ is misleading.

References

- [1] S. W. Hawking and R. Penrose. The singularities of gravitational collapse and cosmology. *Proc. Roy. Soc. Lond. A*, 314:529–548, 1970.
- [2] S. A. Hayward. Signature change in general relativity. *Class. Quantum Grav.*, 9:1851–1862, 1992.
- [3] G. F. R. Ellis, A. Sumeruk, D. Coule, and C. Hellaby. Change of signature in classical relativity. *Class. Quantum Grav.*, 9:1535–1554, 1992.
- [4] T. Dereli and R. W. Tucker. Signature dynamics in general relativity. *Class. Quantum Grav.*, 10:365–373, 1993.
- [5] S. A. Hayward. Junction conditions for signature change. *Preprint gr-qc/9303034*, 1993.
- [6] M. Kossowski and M. Kriele. Signature type change and absolute time in general relativity. *Class. Quantum Grav.*, 10:1157–1164, 1993.
- [7] M. Kossowski and M. Kriele. The Einstein equation for signature type changing spacetimes. *Proc. Roy. Soc. Lond. A*, 446:115–126, 1994.
- [8] M. Kossowski and M. Kriele. Transverse, type changing, pseudo Riemannian metrics and the extendability of geodesics. *Proc. Roy. Soc. Lond. A*, 444:297–306, 1994.
- [9] R. Kerner and J. Martin. Change of signature and topology in a five-dimensional cosmological model. *Class. Quantum Grav.*, 10:2111–2122, 1993.
- [10] J. Martin. Hamiltonian quantization of General Relativity with the change of signature. *Phys. Review D*, 49:5086–5095, 1994.
- [11] C. Hellaby and T. Dray. Failure of standard conservation laws at a classical change of signature. *Phys. Review D*, 49:5096–5104, 1994.
- [12] M. Kossowski and M. Kriele. Smooth and discontinuous signature type change in general relativity. *Class. Quantum Grav.*, 10:2363–2371, 1993.
- [13] J. B. Hartle and S. W. Hawking. Wave function of the universe. *Phys. Review D*, 28(12):2960–2975, 1983.

- [14] M Carfora and G. Ellis. The geometry of classical change of signature. *Preprint*, 1994.
- [15] S. A. Hayward. On cosmological isotropy, quantum cosmology and the weyl curvature hypothesis. *Class. Quantum Grav.*, 10:L7–L11, 1993.
- [16] S. A. Hayward. General laws of black-hole dynamics. *Phys. Review D*, 49:6467–6474, 1994.
- [17] M. Kossowski and M. Kriele. Transverse, type changing, pseudo Riemannian metrics with smooth curvature. *Preprint FI93-CT11, The Fields Institute for Research in Mathematical Sciences*, 1993.
- [18] M. Kossowski. Fold singularities in pseudo Riemannian geodesic tubes. *Proc. Amer. Math. Soc.*, 95(3):463–469, 1985.
- [19] P. R. Garabedian. *Partial differential equations*. Chelsea Publishing Company, New York, 2nd edition, 1986.